

## GENERATION OF VIRTUAL COMBINATORIAL LIBRARIES OF COMPOUNDS

### FIELD OF THE INVENTION

INS A17  
5

The present invention is directed to methods for the generation of virtual combinatorial libraries of small molecules and other ligands. The members or molecules of the combinatorial libraries are generated *in silico*, and are designed to bind to identified target molecules *in silico*. The present invention also includes methods for docking the library members to desired target molecules whereby the library members are bound to such targets *in silico*.

### BACKGROUND OF THE INVENTION

Combinatorial chemistry is a recent addition to the toolbox of chemists and represents a field of chemistry dealing with the synthesis of a large number of chemical entities. This is generally achieved by condensing a small number of reagents together in all combinations defined by a given reaction sequence. Advances in this area of chemistry include the use of chemical software tools and advanced computer hardware which has made it possible to consider possibilities for synthesis in orders of magnitude greater than the actual synthesis of the library compounds. The concept of "virtual library" is used to indicate a collection of candidate structures that would theoretically result from a combinatorial synthesis involving reactions of interest and reagents to effect those reactions. It is from this virtual library that compounds are selected to be actually synthesized.

Project Library (MDL Information Systems, Inc., San Leandro, CA) is said to be a

desktop software system which supports combinatorial research efforts. (*Practical Guide to Combinatorial Chemistry*, A. W. Czarnik and S. H. DeWitt, eds., 1997, ACS, Washington, D.C.) The software is said to include an information-management module for the representation and search of building blocks, individual molecules, complete combinatorial  
5 libraries, and mixtures of molecules, and other modules for computational support for tracking mixture and discrete-compound libraries.

Molecular Diversity Manager (Tripos, Inc., St. Louis, MO) is said to be a suite of software modules for the creation, selection, and management of compound libraries. (*Practical Guide to Combinatorial Chemistry*, A. W. Czarnik and S. H. DeWitt, eds., 1997,  
10 ACS, Washington, D.C.) The LEGION and SELECTOR modules are said to be useful in creating libraries and characterizing molecules in terms of both 2-dimensional and 3-dimensional structural fingerprints, substituent parameters, topological indices, and physicochemical parameters.

Afferent Systems (San Francisco, CA) is said to offer combinatorial library software  
15 that creates virtual molecules for a database. It is said to do this by virtually reacting precursor molecules and selecting those that could be actually synthesized (Wilson, *C&EN*, April 27, 1998, p.32).

While only Project Library and Molecular Diversity Manager are available commercially, these products do not provide facilities to efficiently track reagents and  
20 synthesis conditions employed for the introduction of fragments into the desired compounds being generated. Further, these products are unable to track mixtures of compounds that are generated by the introduction of multiple fragments by the use of multiple reagents. Therefore, it is desirable to have available methods for handling mixtures of compounds, as well as methods for the tracking of chemical reactions or transformations utilized in the  
25 synthesis of individual compounds and mixtures thereof.

## SUMMARY OF THE INVENTION

In accordance with the present invention, there are provided methods for the generation of virtual combinatorial libraries of small molecules. These library molecules or members are generated *in silico*. Library members of larger molecular weight, such as those  
30 that are polymeric in nature, may also be generated using the methods of the present

invention.

The present invention further provides methods for tracking and maintaining in databases, the fragments, reagents and unique combinations of these used for the *in silico* generation of the library members. Methods for interfacing the information necessary for the generation of libraries *in silico*, as instructions designed to direct the actual synthesis of the library members on an instrument such as a parallel array synthesizer, are also provided in the present invention.

The present invention also provides methods for the *in silico* docking of the library members to identified target molecules. According to these methods, individual library members are allowed to bind to the desired target molecule in order to identify those library members that demonstrate high affinity binding to the targets.

While there are a number of ways to identify molecular interaction sites, identify compounds likely to interact with molecular interaction sites of RNA and other biological molecules, synthesize such compounds and analyze their binding, preferred methodologies are described in U.S. patent applications filed on even date herewith and assigned to the assignee of this invention. These application bear U.S. Serial Nos. (Unknown) and have been assigned attorney docket numbers IBIS-0002, IBIS 0004, IBIS-0005, IBIS-0006 and IBIS-0007. All of the foregoing applications are incorporated by reference herein in their entirety.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a compound, compound CI, dissected into its constituent fragments; Figure 2 shows the various identifying characteristics of the fragments comprising compound CI;

Figure 3 shows the various identifying characteristics of the reagents used to introduce the corresponding fragments comprising compound CI;

Figure 4 is a list of transformations that link the fragments and reagents associated with the generation of compound CI;

Figure 5 is a schematic for the introduction of a common fragment using two different reagents;

Figure 6a is a schematic for the use of a single reagent for the introduction of two different fragments into a compound;

Figure 6b is a schematic showing the use of a common reagent for the introduction of a common fragment into the compound which can further be converted into two different fragments within the compound generated;

Figure 7 shows the symbolic addition of fragments yielding a symbolic compound,  
5 compound CI';

Figure 8 is a symbolic reagent table;

Figure 9 is a symbolic fragment table;

Figure 10 is a symbolic transformation table;

Figure 11 shows the generation of individual compounds, compounds C1 and C4, and  
10 a mixture, mixture M1;

Figure 12 shows the generation of further mixture, mixture M2;

Figure 13 shows the generation of an additional mixture, mixture M3;

Figures 14a and 14b show the generation of an additional mixture, mixture M4;

Figure 15 shows tables for tracking compound C1 by the fragments added and or  
15 transformations performed;

Figure 16 shows tables for tracking mixture M1 by the transformations performed;

Figure 17 shows tables for tracking mixture M2 by the transformations performed; and

Figure 18 shows tables for tracking mixture M3 by the transformations performed.

The present invention is directed to computational methods employed for the *in silico*  
20 design and synthesis of combinatorial libraries of small molecules. The library members are generated *in silico*. The present invention also encompasses methods for tracking and storing the information generated during the *in silico* creation of library members into relational databases for later access and use. For the purposes of this specification, *in silico* refers to the creation in a computer memory, i.e., on a silicon or other like chip. Stated otherwise *in silico*  
25 means "virtual."

According to the methods of the present invention, each compound or library member is dissected into its component or constituent parts referred to as fragments. Thus each compound that is generated is considered to be comprised of constituent fragments such that the sum of the molecular formulas of each of the fragments when added together totals the  
30 molecular formula of the compound generated. This dissection can be done in a variety of ways using chemical intuition. Thus a variety of components of fragments may be identified,

each of which lend themselves to readily available reagents or reactions to generate diverse compounds. Further, each fragment is associated with at least one reagent, which represents the necessary chemical to be used to introduce that desired fragment into the compound being generated *in silico*. Dissection of compounds is based on the ease of synthesis of the reagents, commercial availability of the reagents, or a combination of both. Each of the fragments and reagents are stored in a relational database and are described in terms of identifying characteristics in the database. A fragment may be available from a variety of starting materials or reaction schemes. So when a library is being generated, which entails building a database, the fragments used in building that library can be stored in the database using the corresponding set of reagents and reaction conditions. When another library is to be generated, the fragment information stored in the database is now available for use in the generation of the new library of compounds. Similarly, when a third library is being generated, an even greater quantity of fragment, reagent, and reaction information is available in the database. Thus the methods of the present invention represent a dynamic method of building a database associated with building libraries of compounds. Initial library generation requires database input for fragments, reagents and transformations necessary for desired library. As the database grows, however, an increasing number of fragments and reagents are available in the database, which simplifies the generation of subsequent libraries of compounds and makes for more routine combinatorial synthetic efforts which can be accomplished with increasing ease and efficacy.

Fragments that are recorded in the database may be defined using identifying characteristics. Identifying characteristics defining fragments include a structural representation (as a 2-dimensional or 3-dimensional file), name, molecular weight, molecular formula, and attachment points or nodes (which denote sites of attachment or linkage of the fragment to other fragments of the compound being generated *in silico*). For the purpose of describing this invention, 2-dimensional representations are used, which are further simplified by the use of symbolic representations without reference to any particular chemical entities. The symbolic representations as used herein merely shows how fragments can be tracked to further the methods of the present invention. Other identifying characteristics may also be added to the database. Any characteristic that is desired to be tracked may be included in the database, including biological data, chemical reactivity rates, or other physical or chemical

properties. Further, a fragment may also be created by modifying a reagent, and such modifications can be added to the database in terms of changes made to the reagent structure. Some of the identifying characteristics associated with any fragment may be common to those of the corresponding reagent. The related fragment thus created can then be stored in the relational database.

Identifying characteristics defining reagents include a structural representation, name, molecular weight, molecular formula, and source, such as a commercial source or a unique compound defined by the user. In case of a commercial source for the reagent, a catalog number or a link to a web page can be provided. Some commonalities may exist between the identifying characteristics associated with a reagent and those associated with the related fragment.

Further, in accordance with the present invention, a compound is the sum of various transformations. Transformation is the nomenclature attributed according to the present invention to a chemical synthesis. A transformation is a 1:1 link between a fragment and a reagent. Thus each transformation describes a unique conversion of a reagent into the corresponding fragment as introduced into a compound. When the compound being generated *in silico* is broken down into its component fragments, and the corresponding reagents have been identified, each fragment is linked to the corresponding reagent in a 1:1 relationship in order to describe a transformation. Thus, according to the present invention, a transformation may be viewed as the source of a fragment, thereby linking that fragment to a particular synthetic method or reaction. This description of a transformation according to the methods of the present invention also includes any auxiliary reagents or conditions used to effect the reaction denoted by the transformation, such as temperature and pressure requirements, catalysts, activators, solvents, or other additives.

Each combination of a fragment and reagent in a 1:1 link comprises a different transformation. Therefore, each transformation is unique. The present invention allows the tracking of fragments in terms of the reaction or transformation in which those fragments are introduced into the compounds of the library. Thus the database describes not only the compounds generated in terms of their constituent fragments, but also in terms of the synthetic pathways to produce those compounds, *i.e.* the related transformations to generate the library compounds. In this manner, a user of the present invention can generate a virtual library of

compounds by simply selecting the fragments desired. Alternately, a user can also generate the compounds by selecting the chemical pathways required for actual synthesis of the compounds. This is accomplished by selecting the appropriate transformation associated with the generation of the desired compounds. Here, the user uses intuition or an *in silico* expert system to assist in selecting those transformations that are expected to allow generation or synthesis of the desired compounds. Each of the transformations created *in silico* is stored in the relational database and described in terms of identifying characteristics. Identifying characteristics defining transformations include the fragment, the reagent, and any auxiliary reagent or conditions necessary to effect the conversion of the reagent into the fragment as incorporated into the compound.

For example, consider in Figure 1 the *in silico* generation of compound CI according to the methods of the present invention. As shown in Figure 1, upon dissection of CI (molecular formula of  $C_{12}H_{18}N_2O_5S_1$ ), its constituent fragments can be denoted as  $F_i$  (molecular formula of  $H_2NO$ ),  $F_{ii}$  (molecular formula of  $C_5H_9NO$ ), and  $F_{iii}$  (molecular formula of  $C_7H_7O_3S$ ).  $F_i$  can also be a hydroxyl amine moiety linked to a solid support, *i.e.* P-O-NH, wherein P is a solid support. The sum of the molecular formulas of each of the fragments totals the molecular formula of compound CI.

As shown in Figure 2, each of the fragments,  $F_i$ ,  $F_{ii}$ , and  $F_{iii}$ , are stored in a relational database, and are described in terms of identifying characteristics including a structural representation (which may be 2-dimensional or 3-dimensional), an identifier or name, molecular formula and attachment points or nodes which signify sites on the fragment which are linked to other fragments in compound CI. Other information such as molecular weight can also be associated with the fragment in the database.

As shown in Figure 3, each of the corresponding reagents ( $R_i$ ,  $R_{ii}$ , and  $R_{iii}$ ) are also stored in the relational database, and described in terms of identifying characteristics. Identifying characteristics used to define the reagents include a structural representation, and identifier or name and molecular formula. As with the fragment, other associated information such as molecular weight and source (such as a commercial source verses user-supplied, amount on hand, special handling, etc.) can also be stored in database in association with the individual reagents.

Next, each of the transformations associated with the *in silico* generation of compound

CI are also stored in the relational database. As shown in Figure 4, transformation  $T_i$  links reagent  $R_i$  with fragment  $F_i$ ,  $T_{ii}$  links  $R_{ii}$  with  $F_{ii}$ , and  $T_{iii}$  links  $R_{iii}$  with  $F_{iii}$  in a 1:1 relationship. Also, associated with each transformation is the necessary reaction condition, so that transformation  $T_i$  is associated with reaction condition alpha,  $T_{ii}$  with reaction condition beta, and  $T_{iii}$  with reaction condition gamma. In the case of transformation  $T_{iii}$ , reagent  $R_{iii}$  may be a hydroxyl amine attached to a solid support so that fragment  $F_{iii}$  can be represented as a hydroxyl amine moiety attached to a solid support.

While each fragment may be arrived at or generated by a unique corresponding reagent, the present invention also encompasses common fragments that may be generated via two or more reagents, so that two or more transformations can lead to the same fragment. As shown in Figure 5, the common fragment  $\text{CH}_3\text{-CH}_2\text{-C(=O)-}$  may be arrived at via transformation A, which employs reagent X (an acid chloride),  $\text{CH}_3\text{-CH}_2\text{-C(=O)Cl}$ . The common fragment can also be introduced into a compound being generated *in silico* via transformation B, which employs reagent Y (an acid anhydride),  $\text{CH}_3\text{-CH}_2\text{-C(=O)-O-C(=O)-CH}_2\text{-CH}_3$ . Therefore, in accordance with the methods of the present invention, a common fragment can be introduced into the compound via two or more different reagents, and thus via two or more distinct transformations.

Alternately, a common reagent may be employed to effect two or more conversions forming two or more different fragments. This then represents two or more different transformations associated with different conditions. For example, as shown in Figure 6a, common reagent Z,  $\text{CH}_3\text{-CH}_2\text{-NH}_2$ , can be employed to introduce an alkene fragment into the compound under conditions favoring Schiff's base formation. This represents transformation X. The same common reagent Z, however, can also be employed to introduce an amide fragment into the compound by using a different set of conditions, constituting transformation Y. Thus, a common reagent can introduce two or more different fragments into final compounds being generated *in silico*, and can be associated with two or more transformations depending upon the conditions associated with each of those transformations.

Additionally, once a fragment has been introduced into a compound, it can be further modified and converted into yet another fragment without effecting any other chemical changes within the compound formed. As an example, shown in Figure 6b, consider common reagent Z',  $\text{CH}_3\text{-CH}_2\text{-C(=O)CH}_2\text{-Cl}$ . Common reagent Z' corresponds to a fragment having



the structure  $\text{CH}_3\text{-CH}_2\text{-C(=O)CH}_2\text{-}$ . Common reagent Z' may be used to introduce an alkene fragment into the final compound, representing transformation X', under conditions favoring reduction and dehydration. Common reagent Z', however, can also be used to introduce a hydroxyalkyl fragment into the final compound under conditions favoring reduction. This  
5 represents transformation Y'.

The present invention may be described more generally, in terms of symbolic representations. Symbolic representations are used to describe the methods of the present invention because such representations are not limited to any particular chemistry. Symbolic representations merely denote the manner of using the present invention with multiple  
10 chemical entities. Each symbol used in the representations describing the present invention may represent one compound or multiple compounds because the present invention is not limited to tracking a single compound, but may be used to track a vast variety of compounds that can be generated.

Figure 7 shows the symbolic addition of fragments which yields compound C1'. The  
15 fragments have structures  $F_i$ ,  $F_{ii}$ , and  $F_{iii}$  that are added sequentially to yield compound C1'. Structures  $F_i$ ,  $F_{ii}$ , and  $F_{iii}$  are symbolic representations of the fragments that constitute compound C1'. These fragments can be stored in the relational database with the corresponding identifying characteristics for each of them, including the structural representation, name, molecular formula, and attachment sites or nodes. A visual inspection  
20 of compounds C1 and C1' reveals the commonality between the chemical compound C1 and the symbolic representation of a compound C1' as well as the chemical structure of the fragments and the symbolic structure of the fragments.

A symbolic reagent table is shown in Figure 8. Reagents R1 to R10 can be described in terms of their structure, name, molecular formula, molecular weight, and source as well as  
25 other information that might be desired to be associated with the reagents.. R3 and R4 are two different reagents, but may be used to introduce the same fragment into a compound. This depends upon the reaction conditions used as reagent R3 is used in a transformation associated with one set of conditions, while reagent R4 is used in another transformation associated with a different set of conditions. Also, reagent R5 is comprised of a mixture of two reagents or  
30 components. These may be (R)- and (S)-stereoisomers, D- and L-isomers, or may be two completely different reagents. While R5 here is represented as a mixture of only two reagents

or components, it will be recognized by the art-skilled that the methods of the present invention may be practiced using a mixture of two or more reagents. Typical reagent mixtures used in constructing libraries might have four, five or more individual reagent constituting the mixture.

5           Figure 9 shows a symbolic fragment table. Fragments F1 to F8 are stored in the relational database with identifying characteristics that include a structural representation, name, molecular weight, molecular formula, and attachment sites or nodes. This table depicts symbolic representations of the various fragments that are introduced into the compounds of the library by the use of reagents symbolized in Figure 8. Thus it can be seen that fragment  
10 F1 can be introduced into the compound by employing reagent R1. In fragment F1, X is an identifier for an attachment site. This indicates that X is the site at which F1 attaches to another fragment in a compound. Similarly, fragment F2 may be introduced into a compound (attaching at its X site) by employing reagent R2.

          Fragment F3, however, can be introduced into the compound by the use of either  
15 reagent R3 or R4. This allows for selection in the choice of the reagent used, and also allows for the consideration of the compatibility of the chemistries involved in the introduction of other fragments into the compound. Next, fragment F4 (which is a mixture of fragments) can be introduced via the use of reagent R5, which is a mixture of reagents, as shown in Figure 8.

20           Fragment F5 has two attachment sites, indicating that other fragments can attach at sites X and Y when F5 has been incorporated into a compound. The presence of two attachment sites indicates that two attachments may be undertaken to build a compound when dealing with F5. Here again, as before, F5 can be introduced into the compound using either of reagents R6 or R7, depending upon the reaction conditions used and the chemistries  
25 involved when introducing other fragments to build the compound.

          Fragments F7 and F8 can be introduced into a compound being created *in silico* by employing reagents R9 and R10, respectively. Both these fragments have three attachment sites, indicating that three attachments to other fragments can occur when using these fragments to build a compound *in silico*. While fragments F7 and F8 have three attachment  
30 sites, it is recognized by the art-skilled that more than three attachment sites may be present in a fragment, allowing for more attachments to the fragment upon introduction into a

compound (with the use of an appropriate reagent).

With the fragment and reagent tables in place in the relational database, a transformation table is created in accordance with the methods of the present invention, by linking a fragment with a reagent to form a unique transformation. Figure 10 shows a symbolic transformation table where a fragment is linked to a reagent in a 1:1 relationship. The identifying characteristics describing each transformation include a 1:1 link (a one to one link) between a fragment and a reagent, and the reaction conditions which include, solvent, concentration, temperature and pressure requirements, or auxiliary reagents necessary to effect the introduction of the fragment into the compound by using an appropriate reagent. Auxiliary reagents include catalysts, activators, acids, bases or other chemicals or additives necessary to effect the fragment introduction described. For example a base can always be added with an alkyl halide to scavenge the acid generated with use of the alkyl halide.

As seen in Figure 10, transformation T1 links fragment F1 with reagent R1. T1 also specifies the reaction conditions ( $\alpha$ ) associated with this 1:1 link. Similarly, T2 links F2 with R2 under conditions  $\beta$ . Transformations T3 and T4 are each unique transformations despite being associated with a common fragment, F3. Transformation T3 links common fragment F3 with reagent R3 under conditions  $\alpha$ , while transformation T4 links the common fragment F3 with another reagent, R4, under the different conditions, conditions  $\delta$ . For example reagent R3 might be an alkyl chloride while R4 might be an alkyl iodide. While these reagents are similar (they are both alkyl halides), they might be used under different reaction conditions. Use of different reagents to effect the introduction of the same fragment into the compound being generated *in silico* represents two unique transformations. This indicates two distinct or unique synthetic ways of introducing the same fragment into the compound. Depending upon the totality of the chemical steps involved in synthesizing the compound, one transformation may be preferred over other transformations that introduce the same fragment into the compound.

Transformation T5 links fragment F4 with reagent R5. R5 is a mixture of reagents, such as (R)- and (S)-stereoisomers, D- and L-isomers, or two or more different reagents. As a result, use of R5 leads to the introduction of a mixture of fragments F4 into the compound. The art-skilled will recognize that the multiple reagents in R5 are selected such that they are capable of being mixed together, do not react with each other, and react under similar reaction

conditions. For example, R5 may be comprised of a mixture of acid halides. These do not react with each other, but do react similarly with a nucleophile under similar conditions. It is also recognized by the art-skilled that a reagent is not limited to only one or two components or constituent reagents, but in fact may comprise of two, three, four, five or more reagents or components.

When using a mixture of reagents, each of the individual component reagents may have different chemical reactivity rates. If a correction is not made for this, this could result in their products being unequally represented in the product compounds. This is solved by adjusting the concentration of each reagent in the reaction mixture relative to the other reagents in the mixture such that the relative rates are the same. This is effected by comparing to the reactivity of each of the reagents to a chosen standard reagent. The standardized reactivity rates can then be used to adjust the concentration of each constituent reagent in the reagent mixture to compensate for the varied reaction rates. Thus a mixture of reagents with different reaction rates may be used in one reagent mixture to still generate equivalent quantities of the desired compounds in the library.

Transformations T6 and T7 are similar to transformations T3 and T4 except that conditions identifying each of these transformations are different. Transformation T6 links fragment F5 with reagent R6 under conditions  $\epsilon$ , while transformation T7 links the same fragment F5 with a different reagent R7 under different conditions (condition  $\alpha$ ). As the conditions associated with transformations T6 and T7 are different, this allows selection of compatible chemistries with other fragments during any particular synthesis being used. This is a very useful and very important consideration in actually synthesizing real libraries. When it is desired to introduce fragment F5 into the compound, the actual chemistries used to build the compound can be initially be considered in selecting transformation T6 or T7, and thus reagents R6 or R7. This is in direct opposition to any chemical database generator that only considers the compound structure not the actual chemistries necessary to build a compound.

Transformations T9 and T10 link fragment F7 with reagent R9 and fragment F8 with reagent R10, respectively. Both transformations are identified to be associated with reaction conditions  $\gamma$ . Fragments F7 and F8 have three attachment sites, but it is recognized that these fragments may have more than three attachment sites, thereby increasing the complexity of the compounds generated, and increasing the number of rounds that may be employed to

attach other fragments. For the three sites illustrated, if three sets of different reagent mixtures each have five reagents in the set are used, then 125 compounds will be generated for fragment F7 and a further 125 compounds will be generated for fragment F8.

The methods of the present invention may be used to generate single compounds or mixtures of compounds. A mixture comprises two or more compounds and may involve the use of two or more reagents (thus introduction of two or more fragments) at the outset of library generation, introduction of a mixture of reagents (thus a mixture of fragments) at a subsequent stage of library generation, or a combination of both such techniques. Figures 11 and 12 illustrate this aspect of the present invention.

As shown in Figure 11, the methods of the present invention may be used to generate single compounds such as C1 and C4, or may also be used to generate a mixture of compounds, M1, comprising compounds C2 and C3. Library generation commences with selecting fragment F7 (with three attachment sites), in the first round (*i.e.* round *n*). In the second synthesis round (*i.e.* round *n*+1), F7 is combined with fragment F2, constituting synthetic pathway P1a, and resulting in the formation of complex fragment CF1. F7 possesses three attachment sites (*i.e.* X, Y and Z). Thus round *n*+1 will not be complete until each of X, Y and Z have been used, if desired, to attach other fragments to. Stepping around each of X, Y and Z, and attaching fragments to these sites, occurs in that sequential order. Once sites X, Y and Z of the fragment selected in the first synthesis round (*i.e.* round *n*) have been exhausted, stepping around the attachment sites present in the next added fragment constitutes the next synthesis round (*i.e.* the third synthesis round, or round *n*+2). Here again, when all desired attachment sites on this fragment have been used, that particular synthesis round is complete. This attachment iteration around the desired and available attachment sites of the fragments added continues until the desired compounds have been generated.

As shown in Figure 11, CF1 is next subjected to synthetic pathway P1b wherein fragment F1 is introduced into CF1, thereby forming complex fragment CF2. CF2 is then subjected to synthetic pathway P1c wherein fragment F5 is added to CF2, leading to the formation of complex fragment CF3. This completes synthesis round *n*+1 (*i.e.* the second round of fragment introduction, or synthesis, to build the compound). As fragment F5 has two attachment sites, CF3 has an available attachment site (*i.e.* site Y). Introduction of fragments to this site (Y site) constitutes synthesis round *n*+2 (*i.e.* the third round) because all the desired

attachment sites on the previously added fragment have been exhausted. Next, CF3 is subjected to synthetic pathway P2 wherein fragment F4 is introduced into CF3 at attachment site Y. As F4 is a mixture of two components, a mixture (M1) of two compounds, C2 and C3, is generated.

5 A single compound, however, may also be generated using the present scheme of fragment introduction. Thus, compound C1 can be generated by subjecting CF3 to synthetic pathway P1d wherein CF3 is combined with fragment F3, which attaches to site Y in CF3. The introduction of fragment F3 into CF3 constitutes the third synthesis round (*i.e.* round  $n+2$ ), leading to the generation of C1.

10 Alternately, CF3 can be subjected to synthetic pathway P3a wherein fragment F6 is introduced into CF3 to form CF4. This represents the third synthesis round (*i.e.* round  $n+2$ ). CF4 has one more available attachment site (*i.e.* site Y) to which fragment F2 may be attached via synthetic pathway P3b. This leads to the generation of compound C4 which is a compound of increased complexity because of the number of attachment sites on the chosen  
15 fragments and synthetic pathways employed. The addition of fragment F6 to CF4 constitutes the third synthesis round (*i.e.* round  $n+2$ ). Addition of fragment F2 to CF4 represents the fourth synthesis round, or round  $n+3$ , because P3b involves addition of a fragment (fragment F2) onto a site (*i.e.* site Y in CF4) which has been generated by adding fragment F6 to CF3, thus exhausting the available attachment sites on the previously added fragment in CF4 (*i.e.*  
20 fragment F5). That is, the addition of fragment F6 completed round  $n+2$  (or the third synthesis round) because F6 attached to the last available attachment site on CF3 (*i.e.* site Y in CF3).

For the reactions effected at path P1c in Figure 11, a single fragment (F5) can be added to CF2 via use of either reagents R6 or R7 (as thus via the transformations associated with R6  
25 and R7). While these additions are represented as two unique transformations for the purpose of tracking in the database on the invention, these additions in effect perform the same chemical conversion. Thus, the simultaneous tracking of compounds generated according to the methods of the invention is useful not only in working with virtual libraries of compounds, but also provide the user with a choice of synthetic pathways along which the compounds can  
30 be actually synthesized. This tracking aspect of the present invention is, therefore, a novel and unique way to account for the fragments being introduced, the related transformations (or

reactions) associated with the fragments, and the alternate transformations that lead to the introduction of a common fragment into the desired compounds. The present invention allows not only the tracking of individual compounds that are generated by the use of multiple reagents, but also allows for the simultaneous tracking of multiple compounds that are generated via multiple transformations. While the methods described herein represent the tracking aspects of the invention in terms of symbolic representations or tables, it is recognized by the art-skilled that a variety of computer algorithmic codes and techniques may be employed for the individual or simultaneous tracking aspects described above.

The present invention further provides methods for the one-pot generation of mixtures of compounds by commencing the library generation using different starting fragments in a one-pot fashion. One-pot generation or synthesis of compounds refers to the formation of multiple compounds in a single reaction vessel (*i.e.* one pot). This is possible if compatible chemistries are selected. Examples of such single vessels include but are not limited to multiple well plates, e.g. a 96-well plate, reactions flasks, e.g. a 25 mL flask, or even an industrial reactor. The reactions, or transformations, are performed in one vessel regardless of the size of the reaction vessel. The concept of one-pot synthesis is irrelevant to the generation of virtual libraries of compounds as these virtual libraries are merely generated *in silico*. The concept of one-pot synthesis becomes relevant, however, when the actual synthesis of libraries of compounds is to be undertaken. Thus the compounds can be tracked separately for compound building in order to generate distinct chemical structures, however, they can be group together for synthesis allowing them to be made in the same "pot."

An example of a one-pot synthesis was shown in Figure 11 with the addition of the complex reagent R5 to form mixture M1. A further one-pot synthesis is shown in Figure 12, where a further mixture of compounds is generated. Mixture M2 comprising compounds C1 and C5 can be generated by starting with fragments F7 and F8 in the first synthesis round (*i.e.* round n). Each of these fragments have three attachment sites onto which other fragments can be introduced. As a result, subjecting the two fragments to synthetic pathway P1a wherein F7 and F8 are combined with fragment F5 at site X, results in the one-pot formation of complex fragments CF1 and CF5. CF1 and CF5 are next subjected to synthetic pathway P1b wherein fragment F1 is introduced into CF1 and CF5 at site Y, thereby forming complex fragments CF2 and CF6. CF2 and CF6 are next subjected to synthetic pathway P1c wherein

fragment F5 is introduced into these complex fragments at site Z, forming CF3 and CF7. This completes the second synthetic round (*i.e.* round  $n+1$ ). As fragment F5 contains two attachment sites, after introduction into CF3 and CF7, there is still available an attachment site (*i.e.* site Y) for further introduction of another fragment. Thus CF3 and CF7 are converted to  
5 a mixture (M2) of compounds C1 and C5 via synthetic pathway P1d wherein CF3 and CF7 are combined with fragment F3 which attaches to the Y site on fragment F5 in CF3 and CF7. The introduction of fragment F3 at site Y in CF3 and CF7 represents the third synthetic round (*i.e.* round  $n+2$ ).

Yet another symbolic example of the one-pot generation of mixtures of compounds,  
10 in accordance with the present invention, is shown in Figure 13. *In silico* generation of compounds commences with the selection of fragment F7, which has three sites of attachment (X, Y, and Z). This represents the first synthesis round (*i.e.* round  $n$ ). Next, F7 is subjected to synthetic pathway P1a wherein F7 is combined with fragment F2. F2 attaches to site X on fragment F7, forming complex fragment CF1. At this stage, CF1 is subjected to two synthetic  
15 pathways, P1b and P1b'. P1b employs fragment F1 which is introduced onto site Y on CF1, thereby forming complex fragment CF2, while P1b' employs fragment F3 which is introduced onto site Y on CF1, thereby forming complex fragment CF8. Thus a mixture of complex fragments (CF2 and CF8) are formed. Both fragments, F1 and F3 can be introduced together (such as from a single reagent bottle when actual synthesis is being undertaken) for the one-  
20 pot generation of compounds if the chemistries associated with introduction of these fragments into the compounds are compatible. If not, these fragments can be introduced separately. Next, CF2 and CF8 are subjected to synthetic pathway P1c wherein both complex fragments are combined with fragment F5 which attaches to site Z on CF2 and CF8, thereby forming complex fragments CF3 and CF9. The formation of CF3 and CF9 completes the  
25 second synthesis round (*i.e.* round  $n+1$ ). As fragment F5 has two sites of attachment, site Y is still available for attachment to another fragment. Therefore, CF3 is subjected to synthetic pathway P3 wherein CF3 is combined with fragment F4. Introduction of F4 represents the third synthesis round (*i.e.* round  $n+2$ ). F4 is a mixture of fragments (and introduced by adding a mixture of reagents), as shown in Figure 9. As a result, synthetic pathway P2 leads to the  
30 generation of compounds C2 and C3. Simultaneously, CF9 combines with fragment F4, via synthetic pathway P2', leading to the generation of compounds C7 and C8. Thus mixture M3



is formed comprising compounds C2, C3, C7 and C8.

The present invention also provides methods for the generation of increasingly complex mixtures of compounds. An example is shown in Figures 14a and 14b where mixture M4 is generated and comprises sixteen compounds. The compounds in mixture M4  
5 can be generated by starting with fragments F7 and F8 in the first synthesis round (*i.e.* round *n*). These fragments can then be combined with fragment F2, which is introduced at site X in each of F7 and F8, forming complex fragment CF1 and CF5. Following this, a mixture of fragments F1 and F3 are introduced into CF1 and CF5 at site Y of these complex fragments, leading to the formation of four complex fragments, CF2, CF6, CF8 and CF11. These  
10 complex fragments are next combined with a mixture of fragments F5 and F6. Both F5 and F6 have two attachment sites such that site X on F5 and F6 attaches to site Z on CF2, CF6, CF8 and CF11 forming a mixture of eight complex fragments, CF3, CF7, CF9, CF12, CF13, CF14, CF15 and CF16. This completes the second synthesis round (*i.e.* round *n*+1). As fragments F5 and F6 have two attachment sites, X and Y, the abovementioned eight complex  
15 fragments have one more available attachment site (*i.e.* site Y) onto which another fragment may be introduced. Attachment of a fragment to site Y on these eight complex fragments represents the third synthesis round (*i.e.* round *n*+2). Next, fragment F4 is introduced into CF3, CF7, CF9, CF12, CF13, CF14, CF15 and CF16. As fragment F4 is a mixture of two constituent fragments, sixteen compounds are generated: C2, C3, C7, C8, C9, C10, C11, C12,  
20 C13, C14, C15, C16, C17, C18, C19 and C20. Thus it can be seen that by using multiple fragments in a one-pot fashion and combining with mixtures of fragments, mixtures of compounds of increasing complexity can be generated. The example in Figures 14a and 14b shows sixteen unique compounds being generated as mixture M4 when the library is generated by starting with two fragments. It is recognized by the art-skilled that if the library  
25 generation is commenced with more than two fragments or multiple fragments are added to the same precursor fragment, even more complex mixtures of compounds can be generated.

The present invention also provides methods for keeping track of fragment addition in the various synthesis rounds. This system of accounting is accomplished by tabulation of the synthesis rounds which are correlated with addition of fragments. While for the purposes  
30 of illustration of the invention, a tabulation method of tracking fragment addition is described herein, it will be recognized by the art-skilled that other algorithms, algorithmic codes,

computer readable mediums and various software coding techniques known to those skilled in the computer arts may be used for such tracking. The tables tracking fragment addition can be used to produce structural representations of compounds and create virtual libraries where actual synthesis of the compounds is not desired. Tables tracking transformations, however, can be used to synthesize compounds by selecting the appropriate transformations, and in the case of multiple transformations, selecting the preferable transformations to introduce the required fragment into the compounds being synthesized.

Figure 15 is descriptive of compound C1 in terms of the fragments added in each synthesis round. The first synthesis round (*i.e.* round n) commences with the selection of fragment F7. This is followed by the sequential addition of fragments F2, F1 and F5 in the second synthesis round (*i.e.* round n+1). Finally, compound C1 is generated by the addition of fragment F3 in the third synthesis round (*i.e.* round n+2). The compounds thus generated can be stored as a 2-dimensional virtual library, or may be converted to a 3-dimensional virtual library that can be used for *in silico* docking to desired target molecules.

For the generation of virtual libraries of compounds and for docking the library members onto target molecules, it suffices to add compounds to the relational database in terms of its fragments to track the addition of fragments in the various synthetic rounds. However, when the actual synthesis of desired compounds of a library is to be undertaken, it becomes necessary to specify the actual synthetic steps, reagents, solvents, concentrations, auxiliary compounds needed and other various synthetic factors in order to effect such an actual synthesis of real chemical compounds. Such synthetic steps, reagents, solvents, concentrations and auxiliary compounds are, in fact, incorporated in to the above described transformations. Thus by employing the concept of transformations, the present invention provides methods to track the compounds generated not only in terms of the fragments added but as well as the synthetic parameters necessary for each synthesis round.

Figure 15 also shows the generation of compound C1 in terms of the various transformations employed in the synthesis rounds. Four synthesis pathways lead to the synthesis of compound C1 because of the availability of multiple transformations that can introduce the same fragment into the compound being synthesized. Thus, as seen in Figure 15, selection of fragment F7 constitutes transformation T9 in the first synthesis round (*i.e.* round n). This is followed by the addition of fragment F2 which is achieved by employing

transformation T2. Next, fragment F1 is added via transformation T1. Fragment F5, however, may be added by employing either reagent R6 via transformation T6 along synthesis paths 1 and 3, or reagent R7 via transformation T7 along synthesis paths 2 and 4. Similarly, the final fragment F3 can be added by using either reagent R3 via transformation T3 along synthesis paths 1 and 2, or reagent R4 via transformation T4 along synthesis paths 3 and 4. Thus Figure 15 shows that compound C1 can be actually synthesized via one of four different synthetic schemes which can be tracked or tabulated and accounted for using the methods of the present invention. Each of the four tables is completely descriptive of each of the four synthetic pathways for the preparation of C1. Thus, a user of the present invention has available all the alternate pathways of performing the same reaction (*i.e.* introducing the same fragment), and can select the preferable or most appropriate synthetic route to preparing the desired compounds.

Figure 16 shows a similar transformation tracking table for compounds C2 and C3 in mixture M1. Synthesis of compounds C2 and C3 commences with selection of fragment F7 which represents transformation T9 (step 1 in Figure 16) in the first synthesis round (*i.e.* round n). Next, F7 is combined with fragment F2 via transformation T2 in the second synthesis round (*i.e.* round n+1) (step 2). In the same round, fragment F1, via transformation T1, and fragment F5, via transformation T7 are added sequentially (steps 3 and 4). Finally, fragment F4 is added in the third synthesis round (*i.e.* round n+2). As F4 is a mixture of two constituent fragments (because of two constituent reagents), the table is duplicated at this stage (step 5) to account for the different synthetic ways in which transformation T5 may be accomplished (*i.e.* T5<sup>1</sup> and T5<sup>2</sup>). Step 5 represents compounds C2 and C3. Thus it can be seen that, in accordance with the present invention, whenever there is more than one reagents associated with a particular transformation, the table is duplicated as many times as there are such reagents.

Figure 17 shows a transformation tracking table for compounds C1 and C5 in mixture M3. As the synthesis commences with two fragments, F7 and F8, tracking begins with two parallel tables (step 1 in Figure 17). In the first synthesis round (*i.e.* round n), F7 is selected via transformation T9, while F8 is selected via transformation T10. The second synthesis round (*i.e.* round n+1) commences at step 2 with the introduction of fragment F2 via transformation T2. In step 3, transformation T1 introduces fragment F1 into the compound.

In step 4, transformation T7 introduces fragment F5. This completes the second synthesis round (*i.e.* round  $n+1$ ). Finally, in the third synthesis round (*i.e.* round  $n+2$ ), transformation T4 is used to introduce fragment F3 (at step 5) producing mixture M2 comprising compounds C1 and C5. In this example, the tables are duplicated early in the synthetic scheme because  
5 of the use of a mixture of fragments F7 and F8 at the outset.

The transformation tracking table for compounds C2, C3, C7 and C8 of mixture M3 are shown in Figure 18. The synthesis of these compounds commences with the first synthesis round (*i.e.* round  $n$ ) in which fragment F7 is selected. This represents transformation T9 (shown in step 1 in Figure 18). Step 2 in Figure 18 depicts the second  
10 synthesis round (*i.e.* round  $n+1$ ) and involves the addition of fragment F2 via transformation T2. While steps 1 and 2 involve single transformations each, step 3 involves two different transformations because two different fragments are being introduced into the compounds through the use of two different reagents. Therefore, at step 3 the table is twice duplicated because two different reagents are being employed to introduce two different fragments via  
15 two different transformations. In step 3, transformation T1 is used to introduce fragment F1 while transformation T3 is used to introduce fragment F3. The second synthesis round (*i.e.* round  $n+1$ ) is completed at step 4 with transformation T7 which introduces fragment F5. In the final synthesis round (*i.e.* the third round or round  $n+2$ ), transformation T5 is used to introduce fragment F4. As F4 is a mixture of two constituent fragments, each table at step 5  
20 is twice duplicated for transformations  $T5^1$  and  $T5^2$  which represent each of the constituent fragments of F4.

These figures represent merely one manner in which the various fragments, reagents and transformations may be tracked during the generation or synthesis of single compounds or mixtures of compounds. It will, however, be recognized by the art-skilled that various  
25 other algorithm schemes may be employed to track and account for the fragments being introduced via transformations when compounds are being generated *in silico*.

The library members or compounds generated according to the methods of the present invention can be converted into three-dimensional representations using commercially available software. Next, the compounds, in their three-dimensional structures can be docked  
30 onto identified targets, also represented as three-dimensional structures.

Docking of these library members (or ligands) entails the *in silico* binding of the

members to desired target molecules. A variety of theoretical and computational methods are known in the literature to study and optimize the interactions of small molecules with biological targets such as proteins and nucleic acids. These structure-based drug design tools have been very useful in modeling the interactions of proteins with small molecule ligands and in optimizing these interactions. Typically this type of study was performed when the structure of the protein receptor was known by querying individual small molecules, one at a time, against this receptor. Usually these small molecules had either been co-crystallized with the receptor, were related to other molecules that had been co-crystallized or were molecules for which some body of knowledge existed concerning their interactions with the receptor. A significant advance in this area was the development of a software program called DOCK that allows structure-based database searches to find and identify the interactions of known molecules to a receptor of interest (Kuntz *et al.*, *Acc. Chem. Res.*, **1994**, 27, 117; Gschwend and Kuntz, *J. Comput.-Aided Mol. Des.*, **1996**, 10, 123). DOCK allows the screening of molecules, whose 3D structures have been generated *in silico*, but for which no prior knowledge of interactions with the receptor is available. DOCK, therefore, provides a tool to assist in discovering new ligands to a receptor of interest. DOCK can thus be used for docking the compounds prepared according to the methods of the present invention to desired target molecules.

The DOCK program has been applied to protein targets and the identification of ligands that bind to them. The DOCK software program consists of several modules, including SPHGEN (Kuntz *et al.*, *J. Mol. Biol.*, **1982**, 161, 269) and CHEMGRID (Meng *et al.*, *J. Comput. Chem.*, **1992**, 13, 505). SPHGEN generates clusters of overlapping spheres that describe the solvent-accessible surface of the binding pocket within the target receptor. Each cluster represents a possible binding site for small molecules. CHEMGRID precalculates and stores in a grid file the information necessary for force field scoring of the interactions between binding molecule and target. The scoring function approximates molecular mechanics interaction energies and consists of van der Waals and electrostatic components. DOCK uses the selected cluster of spheres to orient ligands molecules in the targeted site on the receptor. Each molecule within a previously generated 3D database is tested in thousands of orientations within the site, and each orientation is evaluated by the scoring function. Only that orientation with the best score for each compound so screened is

stored in the output file. Finally, all compounds of the database are ranked in order of their scores and a collection of the best candidates may then be screened experimentally.

Using DOCK, ligands have been identified for certain protein targets. Recent efforts in this area have resulted in reports of the use of DOCK to identify and design small molecule  
5 ligands that exhibit binding specificity for nucleic acids such as RNA double helices. While RNA plays a significant role in many diseases such as AIDS, viral and bacterial infections, few studies have been made on small molecules capable of specific RNA binding. Compounds possessing specificity for the RNA double helix, based on the unique geometry of its deep major groove, were identified using the DOCK methodology (Chen *et al.*,  
10 *Biochemistry*, **1997**, *36*, 11402; Kuntz *et al.*, *Acc. Chem. Res.*, **1994**, *27*, 117). Using a recent X-ray structure for r(UAAGGAGGUGAU).r(AUCACCUCUUA) as the model structure for the A-form RNA duplex, DOCK identified several aminoglycosides as candidate ligands, characterized by shape complementarity to the RNA groove. Binding experiments then revealed that one of these aminoglycosides not only bound preferentially to RNA over B-form  
15 DNA but also that the ligand binds in the targeted RNA major groove. Recently, the application of DOCK to the problem of ligand recognition in DNA quadruplexes has also been reported (Chen *et al.*, *Proc. Natl. Acad. Sci.*, **1996**, *93*, 2635).

As yet there has been no report of the evaluation of virtual libraries against RNA targets. Certain reports of the generation of virtual libraries are available from the standpoint  
20 of library design, generation, and screening against protein targets. Likewise, some efforts in the area of generating RNA models have been reported in the literature. However, there are no reports on the use of structure-based design approaches to query virtual libraries against three-dimensional models of RNA structure so as to identify ligands, such as small molecules, oligonucleotides or other nucleic acids, that bind to such targets. The present invention  
25 provides a solution to this problem by allowing the building of three-dimensional models of RNA structure, the building of virtual libraries of ligands, including small molecules, polymeric compounds, oligonucleotides and other nucleic acids, screening of such virtual libraries against RNA targets *in silico*, scoring and identifying the best potential binders from such libraries, and finally, synthesizing such molecules in a combinatorial fashion and testing  
30 them experimentally to identify new ligands for such targets.

The methods of the present invention aid in the drug discovery process by allowing

the identification of those library members which bind with high affinity to the target molecules and, therefore, represent molecules that may be actually synthesized and developed as lead drug candidates.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217